


Zika in Montana

2017 Summary



Confirmed Zika Cases Reported in Montana



In 2017, Montana reported one case of travel associated Zika in a non-pregnant female who traveled to a Zika affected area.

All cases reported in Montana have been associated with travel.



Gender



27

MEN TESTED




84

WOMEN TESTED

Tests conducted at the Montana Public Health Laboratory

111 People Tested

Type of Test



● Serum PCR ● Urine PCR ● IgM


Common Symptoms Associated with Zika



Conjunctivitis



Headache



Rash



Muscle Pain



Joint Pain



Fever

Many people infected with Zika will not have symptoms. Those that are symptomatic often experience mild clinical illness.

Prevention Recommendations for Zika

Before Traveling

- Know before you go. Are you traveling to a Zika affected area?
- Pregnant women should avoid travel to a Zika affected area.
- Contact your health care provider or local health department if you have questions or concerns.



While Traveling

- Wear long sleeves and pants.
- Use repellent with DEET.
- Use condoms as Zika can be sexually transmitted.



After Traveling

- Continue using insect repellent for 3 weeks after you return from your trip.
- Protect your partner. If you're a woman, use condoms or do not have sex for 2 months. If you're a man, use condoms or do not have sex for 6 months.
- Contact your health care provider or local health department if you have questions or concerns.

